

# DOMINANT ENERGY CONDITION AND CAUSALITY FOR SKYRME-LIKE GENERALIZATIONS OF THE WAVE-MAP EQUATION

WILLIE WAI-YEUNG WONG

ABSTRACT. It is shown in this note that a class of Lagrangian field theories closely related to the wave-map equation and the Skyrme model obeys the dominant energy condition, and hence by Hawking's theorem satisfies finite speed of propagation. The subject matter is a generalization of a recent result of Gibbons.

## 1. INTRODUCTION

Recently Gibbons showed [Gib03] that the Skyrme model obeys the dominant energy condition, and thus settling the problem of causality for that equation. In this note we will give a different proof of the same fact that easily generalizes to a class of Lagrangian field theories that includes, as special cases, the wave-map equation, the Skyrme model, and the Born-Infeld model.

Let  $(M, g)$  be an  $m+1$  dimensional Lorentzian manifold, where sign convention is taken to be  $(-, +, +, \dots)$ , and let  $(N, h)$  be an  $n$  dimensional Riemannian manifold. Let  $\phi : M \rightarrow N$  be a  $C^1$  map. Then the action of  $\phi$  can be used to pull back the metric  $h$  onto  $M$  as a positive semi-definite quadratic form on  $TM$ , we write it as

$$\phi^*h(X, Y) = h(d\phi \cdot X, d\phi \cdot Y)$$

where the left hand side is evaluated at a point  $p \in M$  and the right hand side at the point  $\phi(p) \in N$  for  $X, Y \in T_pM$ . Composing with the inverse metric  $g^{-1}$  we obtain the so-called *strain tensor*  $D^\phi$ , a section of  $T_1^1M$ :

$$(1) \quad D^\phi = g^{-1} \circ \phi^*h ,$$

thus at every point  $p$ ,  $D^\phi$  is a linear transformation of  $T_pM$ . Now, if  $g$  were a Riemannian metric, then for a fixed basis of  $T_pM$ , the matrix  $(D^\phi)$  is positive semi-definite. This is, unfortunately, no longer true in the Lorentzian case, and thus the eigenvalues of  $(D^\phi)$  are in general complex.

Let  $\{\lambda_1, \dots, \lambda_k\}$  denote the non-zero eigenvalues, counted with multiplicity, of  $(D^\phi)$ . Note that by elementary linear algebra, using that  $g$  is non-degenerate and  $h$  is positive definite, one easily sees that

$$(2) \quad k \leq \text{rank}(d\phi) \leq \min(m+1, n) .$$

---

2000 *Mathematics Subject Classification.* 53Z05.

The author is supported by the Commission of the European Communities, ERC Grant Agreement No 208007.

Recall the elementary symmetric polynomials  $s_j(\{\lambda_1, \dots, \lambda_k\})$  given by

$$(3) \quad s_j(\{\lambda_1, \dots, \lambda_k\}) = \sum_{1 \leq \alpha_1 < \alpha_2 < \dots < \alpha_j \leq k} \prod_{i=1}^j \lambda_{\alpha_i}$$

with  $s_0 = 1$  and  $s_j = 0$  for all  $j > k$ . Observe that for the  $(m+1) \times (m+1)$  matrix  $(D^\phi)$ , the elementary symmetric polynomials correspond to the coefficients of the characteristic polynomial, and specifically  $s_1 = \text{tr}(D^\phi)$  and  $s_{m+1} = \det(D^\phi)$ . By abuse of notation, we will write  $s_j(D^\phi)$  when we mean the symmetric polynomials on the eigenvalues of  $(D^\phi)$ . Note that  $s_j(D^\phi)$  is independent of a basis chosen for the vector space  $T_p M$ .

For a given class  $\mathcal{A}$  of maps  $\phi : M \rightarrow N$ , we write

$$\mathcal{U}_{\mathcal{A}} := \{v \in \mathbb{R}^{m+1} \mid v = (s_1, \dots, s_{m+1})(D^\phi), \phi \in \mathcal{A}\}.$$

**Definition 1.** For a given class  $\mathcal{A}$ , let  $\mathcal{U}_{\mathcal{A}} \subset \mathbb{R}^{m+1}$  be an open set that contains  $\mathcal{U}_{\mathcal{A}} \cup \{0\}$ . An admissible function  $F : \mathcal{U}_{\mathcal{A}} \rightarrow \mathbb{R}$  for the class  $\mathcal{A}$  is a sub-additive, concave function, that is  $C^1$  on the interior of  $\mathcal{U}_{\mathcal{A}}$  and continuous up to the boundary.

**Remark 2.** In the definitions above, it only suffices to include terms up to  $s_{m+1}$  in view of (2). Also, observe that sub-additivity and concavity of  $F$  immediately implies that  $F(0) \geq 0$ .

**Definition 3.** A Lagrangian field theory for the class  $\mathcal{A}$  of maps  $\phi : M \rightarrow N$  is said to be a generalized wave-map<sup>1</sup> if the Lagrangian

$$L = F(s_1(D^\phi), s_2(D^\phi), \dots, s_{m+1}(D^\phi))$$

for an admissible  $F$ . Furthermore, we say that the generalized wave-map is defocusing if the first partial derivatives of  $F$  are all non-negative, i.e.  $\partial_i F(v) \geq 0$   $\forall i = 1, \dots, m+1$  and  $\forall v \in \mathcal{U}_{\mathcal{A}}$ . The generalized wave-map is said to be zeroed if  $F(0) = 0$ . Also, we shall refer to a generalized wave-map for which  $\partial_1 F$  is non-vanishing as non-degenerate.

The author hopes that the reason behind the nomenclature will be evident after the proof of the dominant energy condition is developed. We first give some examples of generalized wave-maps:

- Observe that if  $L$  is a linear combination of the symmetric polynomials  $L = \sum c_i s_i(D^\phi)$ , then it is automatically a zeroed generalized wave-map. If in addition the coefficients  $c_i$  are all non-negative, then  $L$  is defocusing. In this case if  $c_1 > 0$  then  $L$  is non-degenerate.
- Take  $(M, g)$  to be a static space-time, i.e.  $M = \mathbb{R} \times \Sigma$  and  $g = -\rho dt^2 \oplus \gamma$  where  $\rho$  is a positive function on  $\Sigma$  and  $\gamma$  is a Riemannian metric on  $\Sigma$ . A static solution to the generalized wave-map is one for which  $\nabla_t \phi = 0$ . The static solution for  $L = s_1$  gives rise to the harmonic map equation from  $\Sigma \rightarrow N$ , while for the case  $n > m$ ,  $L = \sqrt{s_m}$  (recall that  $\dim M = m+1$ ), the equation becomes the minimal surface equation for the embedding of  $\Sigma$  into  $N$ . For the minimal surface equation we take  $\mathcal{U}_{\mathcal{A}} = \mathbb{R}_+^{m+1}$ .
- In the Lorentzian case,  $L = s_1$  is simply the wave-map equation. For  $L = c_1 s_1 + c_2 s_2$  where  $c_1, c_2 > 0$  are coupling constants, we recover the original Skyrme model if we take  $(N, h)$  to be  $SU(2)$  with the bi-invariant metric.

<sup>1</sup>For the lack of a better name. Suggestions are welcome.

In particular, the Skyrme model is a defocusing, zeroed, non-degenerate, generalized wave-map in the terminology adopted in the present paper.

- Let  $b > 0$  be a fixed large constant. We can restrict  $\phi$  to only consider those maps such that the real parts of the eigenvalues of  $D^\phi$  are greater than  $-b$ . Then letting

$$F = \sqrt{\det(b \cdot Id + D^\phi)} - \sqrt{\det(b \cdot Id)}$$

defined on  $\mathcal{U}_A$  being the set where  $\det(b \cdot Id + D^\phi) \geq 0$ , we get the zeroed, defocusing, non-degenerate, generalized wave-map also known as the Born-Infeld model.

Before stating the main theorem, we recall the statement of the dominant energy condition. Recall that the (covariant) stress-energy tensor  $T \in \Gamma(T_2^0 M)$  for a Lagrangian field theory is given by a variational derivative for the Lagrangian *density* relative to the inverse metric,

$$(4) \quad T\sqrt{|\det g|} := \frac{\delta[L\sqrt{|\det g|}]}{\delta g^{-1}} = \left( \frac{\delta L}{\delta g^{-1}} - \frac{1}{2}Lg \right) \sqrt{|\det g|}.$$

**Definition 4.** *The stress-energy tensor  $T$  is said to obey the dominant energy condition at a point  $p \in M$  if  $\forall X \in T_p M$  such that  $g(X, X) < 0$ , the following two conditions are satisfied*

$$(5a) \quad T(X, X) > 0$$

$$(5b) \quad [T \circ g^{-1} \circ T](X, X) \leq 0$$

*unless  $T$  vanishes identically.*

**Remark 5.** *The definition is equivalent to the classical statements (see, e.g. section 4.3 in [HE73] or chapter 9 of [Wal84]) of the dominant energy condition. Observe that (5b) gives that the vector  $g^{-1} \circ T \circ X$  is a causal vector for any time-like vector  $X$ , and (5a) gives that the vector  $g^{-1} \circ T \circ X$  has opposite time-orientation as the time-like vector  $X$ .*

Now we state the main theorem

**Theorem 6.** *A defocusing generalized wave-map obeys the dominant energy condition.*

First we claim that it would suffice to prove the theorem for each  $s_i$ . The following lemma is a general statement on a convexity property of Lagrangian field theories.

**Lemma 7.** *Let  $F$  be a sub-additive, concave function as in Definition 3. Let  $T_i$  denote the stress-energy tensor corresponding to the Lagrangian  $L_i$ . Assume that  $T_i$  obeys the dominant energy condition, or, equivalently, the vectors  $Y_i = g^{-1} \circ T_i \circ X$  are all past-causal for any fixed future time-like  $X$ . Then  $L = F(L_1, \dots, L_{m+1})$  also obeys the dominant energy condition if  $L$  is defocusing.*

*Proof.* The stress-energy tensor  $T$  can be written, using (4), as

$$T = \sum_{i=1}^{m+1} \partial_i F \cdot \frac{\delta L_i}{\delta g^{-1}} - \frac{1}{2} F g = \sum_{i=1}^{m+1} \partial_i F \cdot T_i - \frac{1}{2} (F - \sum_{i=1}^{m+1} \partial_i F \cdot L_i) g.$$

Now considering  $g^{-1} \circ T \circ X$ , the first term in the above expression contributes  $\sum \partial_i F \cdot Y_i$ . Since  $L$  is defocusing, this is a positive linear combination of past-causal vectors, and hence by elementary Minkowskian geometry, is still past-causal. For the second term, since  $g^{-1} \circ g \circ X = X$ , to show that it is also past-causal it suffices to show that

$$F \geq \sum_{i=1}^{m+1} \partial_i F \cdot L_i .$$

But this follows from the fact that  $F$  is concave and  $F(0) \geq 0$ .  $\square$

Unfortunately, it is immediately clear that the theorem may not be strong enough in certain cases for practical application. This is because the vanishing of  $T$  does not guarantee that the map  $\phi$  is trivial. For example, using that  $s_j = 0$  if  $j > \text{rank}(d\phi)$ , it is immediate that if locally around the point  $p$ ,  $\phi$  is one-dimensional, then for any metric  $g$ ,  $s_j(D^\phi) = 0$  if  $j \geq 2$ . On the other hand, this failure of the dominant energy condition arises from a degeneracy which forces the stress-energy tensor to be a null stress tensor in the language of Christodoulou [Chr00], which we can “normalize” away by taking  $L$  to be zeroed. We claim that this is the only possible failure.

**Proposition 8.** *For  $L = s_i$ ,  $T$  obeys the dominant energy condition. Furthermore,  $T = 0$  at a point  $p$  if and only if  $i > \text{rank}(d\phi|_p)$ .*

From this proposition one immediately sees the following energy bound for smooth solutions of the generalized wave-map equation.

**Corollary 9.** *If  $\phi$  is the solution to a defocusing, non-degenerate, zeroed, generalized wave-map, and if  $T = 0$  on a connected open domain  $\mathcal{B}$  of  $M$ , then  $\phi$  is constant on  $\mathcal{B}$ .*

By applying Hawking’s energy conservation theorem (see section 4.3 in [HE73]) the above corollary implies that defocusing, non-degenerate, zeroed, generalized wave-maps have finite speed of propagation (also known as the domain of dependence condition).

In principle, if one has advanced knowledge on a lower bound to the rank of the map  $\phi$ , one can also obtain analogous statements for degenerate cases. We leave such trivial generalizations to the reader.

The author would like to thank Nick Manton and Claude Warnick for introducing him to the problem, and to Mihalis Dafermos for useful discussions.

## 2. A FORMULA FOR THE STRESS-ENERGY TENSOR AND PROOF OF THE MAIN PROPOSITION

In this section, we’ll first derive a formula for the stress-energy tensor. We will begin by making a geometric observation and obtain, almost immediately, a simple tensorial formula for the Lagrangian. Taking the formal variational derivative of the Lagrangian leads to a tensorial expression for the stress-energy tensor, from which Proposition 8 follows via simple linear algebra.

Consider a real vector space  $V$ . Let  $A$  be a linear transformation on  $V$ . Then  $A$  naturally extends to a linear transformation, which we denote  $A^{\sharp j}$ , on  $\Lambda^j(V)$ , the space of alternating  $j$ -vectors over  $V$ . A bit of basic linear algebra (perhaps by extending  $V$  to  $V \otimes_{\mathbb{R}} \mathbb{C}$  and taking a basis of eigenvectors) shows that  $s_j(A)$  is

proportional to  $\text{tr}_{\Lambda^j(V)} A^{\sharp j}$ . Now, letting  $V = T_p M$  and  $A = D^\phi = g^{-1} \circ \phi^* h$ , we observe that

$$(D^\phi)^{\sharp j} = (g^{-1})^{\sharp j} \circ \phi^*(h^{\sharp j}) ,$$

or, to put it in words,  $(D^\phi)^{\sharp j}$  is obtained from first taking the induced metric  $h^{\sharp j}$  on alternating  $j$ -vectors in  $T_{\phi(p)} N$ , pulling it back via  $\phi$ , and composing it with the induced metric  $(g^{-1})^{\sharp j}$  for the alternating  $j$ -forms. In index notation, this can be written as

$$[(D^\phi)^{\sharp j}]_{a_1 \dots a_j}^{b_1 \dots b_j} = g^{b_1 c_1} \dots g^{b_j c_j} (\phi^* h)_{a_1 [c_1] (\phi^* h)_{a_2 [c_2] \dots (\phi^* h)_{a_{j-1} [c_{j-1}] (\phi^* h)_{a_j [c_j]}$$

where the bracket notation in the indices denotes full anti-symmetrization of the  $\{c_1, \dots, c_j\}$  indices. For a Lagrangian proportional to an  $s_j$ , we can assume

$$(6) \quad L = [(D^\phi)^{\sharp j}]_{a_1 \dots a_j}^{b_1 \dots b_j} = g^{a_1 [c_1] \dots g^{a_j [c_j]} (\phi^* h)_{a_1 c_1} \dots (\phi^* h)_{a_j c_j} .$$

It is simple to check, using  $(D^\phi) = \text{diag}(-1, 1, 1, \dots)$  that the above expression has the correct sign: that  $L$  defined thus is a positive multiple of  $s_j$ .

One can also arrive at (6) purely from a linear algebra point of view. Let  $p_j$  be the power sum

$$p_j(\{\lambda_1, \dots, \lambda_k\}) = \sum_{i=1}^k \lambda_i^j .$$

Recall that we have Newton's identity

$$j \cdot s_j = \sum_{i=1}^j (-1)^{i-1} e_{j-i} p_i$$

which allows us to express  $s_j$  as a rational polynomial in  $p_i$ 's. Now, by definition, it is clear that

$$p_j(D^\phi) = \text{tr}[(D^\phi)^j]$$

where  $(D^\phi)^j$  is the  $j$ -fold composition of  $D^\phi$ . It is easy to check then, for some  $E$

$$s_j = g^{a_1 b_1} \dots g^{a_j b_j} E_{b_1 \dots b_j}^{c_1 \dots c_j} (\phi^* h)_{a_1 c_1} \dots (\phi^* h)_{a_j c_j} .$$

Newton's identity reduces to a generating condition for  $E$  based on the Kronecker  $\delta$  symbols,

$$E_b^c = \delta_b^c ,$$

$$j E_{b_1 \dots b_j}^{c_1 \dots c_j} = \sum_{i=1}^j (-1)^{i-1} E_{b_1 \dots b_{j-i}}^{c_1 \dots c_{j-i}} \delta_{b_{j-i+1}}^{c_{j-i}} \delta_{b_{j-i+2}}^{c_{j-i+1}} \dots \delta_{b_j}^{c_{j-i+1}} .$$

A direct computation which we omit here shows that then in fact the invariant  $E_{b_1 \dots b_j}^{c_1 \dots c_j}$  is a positive rational multiple of the generalized Kronecker symbol  $\delta_{b_1 \dots b_k}^{c_1 \dots c_j}$ , from which we recover (6).

Now, the object we are interested in, given a time-like vector  $X$ , is the one-form  $T(X, \cdot)$ . Since  $T$  is tensorial, we can assume  $X$  has unit length. Fix some  $j$ , let the Lagrangian be proportional to  $s_j$  as given by (6). By the symmetry property, we can write  $T(X, \cdot)$  in index notation:

$$(7) \quad T_{ab} X^b = j X^{[b]} g^{a_2 [c_2] \dots g^{a_j [c_j]} (\phi^* h)_{ab} \dots (\phi^* h)_{a_j c_j} - \frac{1}{2} g_{ab} X^b L$$

*Proof of Proposition 8.* Consider a orthonormal basis for  $T_p M$  relative to  $g$ . Since we assumed  $X$  unit, let  $e_0 = X$  and  $\{e_i\}_{1 \leq i \leq m}$  are all space-like. We can take  $j \leq m+1$  as otherwise  $T$  is identically 0. Then we notice that a basis for  $\Lambda^j(T_p M)$  is given by

$$\{e_0 \wedge e_{\alpha_1} \wedge \cdots \wedge e_{\alpha_{j-1}}\}_{1 \leq \alpha_1 < \cdots < \alpha_{j-1} \leq m} \cup \{e_{\alpha_1} \wedge \cdots \wedge e_{\alpha_j}\}_{1 \leq \alpha_1 < \cdots < \alpha_j \leq m} .$$

We write the first set as  $\Lambda_{\perp}^j$  and the second set as  $\Lambda_{\parallel}^j$ . Using the normalization that  $v \wedge w = v \otimes w - w \otimes v$ , we find that each of the element in  $\Lambda_{\perp}^j$  has norm  $-j!$  while the elements in  $\Lambda_{\parallel}^j$  has norm  $j!$ .

To show that  $T(X, X) > 0$  generically, we observe that under the expansion (7), the first term corresponds to

$$\sum_{\omega \in \Lambda_{\perp}^j} \phi^*(h^{\sharp j})(\omega, \omega) ,$$

while the second term corresponds to

$$\frac{1}{2} \left( - \sum_{\omega \in \Lambda_{\perp}^j} \phi^*(h^{\sharp j})(\omega, \omega) + \sum_{\omega \in \Lambda_{\parallel}^j} \phi^*(h^{\sharp j})(\omega, \omega) \right) .$$

So summing them gives

$$\frac{1}{2} \left( \sum_{\omega \in \Lambda_{\perp}^j} \phi^*(h^{\sharp j})(\omega, \omega) + \sum_{\omega \in \Lambda_{\parallel}^j} \phi^*(h^{\sharp j})(\omega, \omega) \right)$$

which is non-negative by the fact that  $\phi^*(h^{\sharp j})$  is a positive semi-definite quadratic form on  $\Lambda^j(T_p M)$ . Furthermore, observe that since  $\Lambda_{\parallel}^j \cup \Lambda_{\perp}^j$  is a basis, its push-forward  $\phi_* \Lambda_{\parallel}^j \cup \phi_* \Lambda_{\perp}^j$  spans  $\Lambda^j(\phi_* T_p^M) \subset \Lambda^j(T_{\phi(p)} N)$ . Thus by the fact that  $h$  (and hence the induced metric  $h^{\sharp j}$ ) is positive definite, we conclude that  $\Lambda^j(\phi_* T_p^M) = \{0\}$ , which proves the assertion that  $T$  vanishes only when  $j > \text{rank}(d\phi)$ .

To show (5b), we observe that

$$X^a T_{ac} g^{cd} T_{db} X^b = -T(X, X)^2 + \sum_{i=1}^m T(X, e_i)^2 .$$

The first thing to note is that  $T(X, e_i)$  does not have any contribution from the second term in (7). For the first term, a quick computation shows that  $T(X, e_i)$  corresponds to

$$\sum_{\eta \in \Lambda_{\parallel}^{j-1}} \phi^*(h^{\sharp j})(e_0 \wedge \eta, e_i \wedge \eta)$$

so

$$\begin{aligned}
\left| \sum_{i=1}^m T(X, e_i)^2 \right| &\leq \left( \sum |T(X, e_i)| \right)^2 \\
&\leq \left( \sum_{i=1}^m \sum_{\eta \in \Lambda_{\parallel}^{j-1}} |\phi^*(h^{\sharp j})(e_0 \wedge \eta, e_i \wedge \eta)| \right)^2 \\
&\leq \frac{1}{4} \left( \sum_{\eta \in \Lambda_{\parallel}^{j-1}} \phi^*(h^{\sharp j})(e_0 \wedge \eta, e_0 \wedge \eta) + \sum_{i=1}^m \phi^*(h^{\sharp j})(e_i \wedge \eta, e_i \wedge \eta) \right)^2 \\
&= \frac{1}{4} \left( \sum_{\eta \in \Lambda_{\parallel}^{j-1}} \sum_{i=0}^m \phi^*(h^{\sharp j})(e_i \wedge \eta, e_i \wedge \eta) \right)^2 \\
&= T(X, X)^2
\end{aligned}$$

And therefore (5b) is satisfied.  $\square$

#### REFERENCES

- [Chr00] Demetrios Christodoulou, *The action principle and partial differential equations*, Annals of Mathematics Studies, Princeton University Press, 2000.
- [Gib03] Gary W. Gibbons, *Causality and the Skyrme model*, Physics Letters B **566** (2003), 171–174.
- [HE73] Stephen William Hawking and George F. R. Ellis, *The large scale structure of space-time*, Cambridge University Press, 1973.
- [Wal84] Robert M. Wald, *General relativity*, University of Chicago Press, 1984.

DEPARTMENT OF PURE MATHEMATICS AND MATHEMATICAL STATISTICS, UNIVERSITY OF CAMBRIDGE, CAMBRIDGE, UK

*E-mail address:* ww278@dpmms.cam.ac.uk